

Name	: XXXXXXXXXXXX	Id	: XXXXXXXXXXXX
Age/Gender	: YY	Vid	: XXXXXXXXXXXX
Referred By	: XXXXXXXXXXXX	Billed	: XXXXXXXXXXXX
		Collected On	: XXXXXXXXXXXX
		Reported On	: XXXXXXXXXXXX

TEST	RESULT	UNITS	Biological Reference Interval
------	--------	-------	-------------------------------

DEPARTMENT OF MOLECULAR BIOLOGY

Orbito Respiratory Bacterial - 1 (1511)
(Method: Multiplex Real Time PCR)

Staphylococcus aureus	NOT DETECTED		
Chlamydia pneumoniae	NOT DETECTED		
Haemophilus influenzae	NOT DETECTED		
Streptococcus pneumoniae	NOT DETECTED		
Moraxella catarrhalis	DETECTED		
Bordetella Spp.	NOT DETECTED		
Haemophilus influenzae (Type B)	NOT DETECTED		
Flu C	NOT DETECTED		

INTERPRETATION

- Staphylococcus aureus:** *Staphylococcus aureus* is a facultative anaerobic Gram-positive coccus, and it is frequently found as a commensal organism in the respiratory tract and on the skin. These bacteria are spread by having direct contact with an infected person, by using a contaminated object, or by inhaling infected droplets dispersed by sneezing or coughing. This versatile bacterium can invade many tissues and then causes a wide spectrum of infections (cutaneous abscesses, endocarditis, septic shock, etc.).
- Streptococcus pneumoniae:** *Streptococcus pneumoniae* remains the leading cause of bacterial meningitis. It is the commonest cause of meningitis between the ages of 1 and 23 months, and above the age of 19. The nasopharynx is the primary site of colonization, and the vast majority of pneumococcal isolates are encapsulated. In the majority of these people, the bacteria is not growing or active and will not cause illness. However, anyone who carries this bacteria can transmit it to others, potentially causing any of the illnesses or pneumococcal meningitis.
- Chlamydia Pneumoniae :** *Chlamydia Pneumoniae* is a type of bacteria that causes respiratory tract infections, such as pneumonia (lung infection). The bacteria cause illness by damaging the lining of the respiratory tract including the throat, windpipe, and lungs. Some people may become infected and have mild or no symptoms. *C. pneumoniae* can also cause lower respiratory tract infections like bronchitis (inflammation or swelling of the airways that carry air to the lungs) and pneumonia (lung infection). Some reports say that people with pneumonia caused by *C. pneumoniae* are more likely to have laryngitis (inflammation of the voice box) compared to people with other types of bacterial pneumonia. It can take 3 to 4 weeks for symptoms to appear after someone has been exposed to the bacteria. Symptoms can also continue for several weeks after they start.
- Haemophilus influenzae B :** *Haemophilus influenzae B* Similar to type A, *Haemophilus influenzae B* is also highly contagious and can have dangerous effects on your health in more severe cases. However, this form can only be spread from human to human.
- Klebsiella pneumoniae:** *Klebsiella pneumoniae* is a Gram-negative, non-motile, encapsulated, lactose-fermenting, facultative anaerobic, rod-shaped bacterium. *Klebsiella* bacteria are mostly spread through person-to-person contact. Less commonly, they are spread by contamination in the environment.
- Moraxella catarrhalis:** *Moraxella catarrhalis* is a gram-negative diplococcus that commonly colonizes the upper respiratory tract. It is a leading cause of otitis media in children, acute exacerbations of chronic obstructive pulmonary disease (COPD), and acute bacterial rhinosinusitis. Symptoms include discolored drainage from the nose, high fever, fatigue, swelling in the face, and pain in the forehead or behind the eyes.
- Bordetella pertussis:** *Bordetella pertussis* is a Gram-negative, aerobic, pathogenic, encapsulated coccobacillus of the genus *Bordetella*, and the causative agent of pertussis or whooping cough. The bacterium is spread by airborne droplets; its incubation period is 7–10 days on average (range 6–20 days). Infection results in colonization and rapid multiplication of the bacteria on the mucous membranes of the respiratory tract. The infection occurs mostly in children under the age of one when they are unimmunized, or



Name : XXXXXXXXXXXX
Age/Gender : YY
Referred By : XXXXXXXXXXXX

Id : XXXXXXXXXXXX
Vid : XXXXXXXXXXXX
Billed : XXXXXXXXXXXX
Collected On : XXXXXXXXXXXX
Reported On : XXXXXXXXXXXX

TEST	RESULT	UNITS	Biological Reference Interval
------	--------	-------	-------------------------------

- *children with faded immunity, normally around the ages 11 through 18. The signs and symptoms are similar to a common cold: runny nose, sneezing, mild cough, and low-grade fever.*

--- End of the Report ---

